

Children from birth to age three years automatically qualify for early intervention services if they have a documented medical diagnosis of any of the following conditions.

NEW: 1p36 Deletion syndrome

49, XXXXY syndrome 9p Deletion syndrome Aicardi syndrome

Albinism Amyoplasia

Angelman syndrome

Aniridia

Anophthalmia Anoxic brain injury Apert syndrome Arthrogryposis

Autism Spectrum Disorders Bardet-Biedl syndrome Brachial Plexopathy

Brain Malformation, congenital

Cerebral Palsy CHARGE syndrome

Cornelia de Lange syndrome Cortical Visual Impairment

Cri du chat syndrome

Cytomegalovirus (CMV), congenital

Deafblind delayed

Delayed Visual Maturation (DVM)

DiGeorge/Velocardiofacial syndrome

Down syndrome (Trisomy 21)

Encephalocele Failure to Thrive

Familial Exudative Vitreoretinopathy

(FEVR)

Fetal Alcohol syndrome

Fragile X syndrome

Goldenhar syndrome

Hearing loss

Herpes, congenital Hunter syndrome

Hurler-Scheie syndrome

Hypoxic Ischemic Encephalopathy

(HIE)

Kleefstra (9q34 Deletion) syndrome

Klinefelter syndrome

Lead Poisoning, venous blood level

≥10mcg/dL

Leber's Congenital Amaurosis

Limb Reduction Defect

Microcephaly Microphthalmia Möbius Sequence Muscular Dystrophy

Muscular Dystrophy, Duchenne

Type Myopathies

Neonatal Abstinence syndrome

(NAS)

Neurofibromatosis Type 1 Neurotrauma, inflicted Niemann-Pick disease

Optic Atrophy

Optic Nerve Hypoplasia Osteogenesis imperfecta

Pervasive Developmental Disorder (PDD), NOS

Phthisis Bulbi

Pierre-Robin syndrome Prader-Willi syndrome

Retinal Detachment

Retinopathy of Prematurity (ROP), Stage 4/5

Rubenstein-Taybi syndrome

Septo-Optic Dysplasia Spina Bifida

Spinal Cord Injury with Cord

Involvement

Spinal Muscular Atrophy (SMA)

Sturge-Weber syndrome

Syphilis, congenital

Tethered Cord syndrome (TCS)

Treacher Collins syndrome

Trisomy 13 (Patau syndrome)

Trisomy 18 (Edwards syndrome) Trisomy 21 (Down syndrome)

Tuberous Sclerosis

Unbalanced Chromosomal

Translocation
VACTERL association
Ventilator dependence
Williams syndrome

Wolf-Hirschhorn syndrome

NICU grads with the following conditions are automatically eligible until their first birthday.

- NEW: NICU Other
- Extremely Low Birth Weight (<1000g)
- Extremely preterm (<28 wks)
- Very low birth weight (<1500g) AND very preterm (28-31 wks)
- Fluctuating tone
- Full/partial NG/NJ/G tube required
- Inconsolability
- Learning to eat difficult/slow
- Poor coord suck/swallow/breath
- Severe sleep disorder
- Significant irritability, neuro-based
- Significant tremors at rest
- Unable to come to quiet alert state
- Unable to take 100% oral nutrition
- Unusually high or low tone

NOTE: When an initial diagnosis is acute in nature, it may be appropriate to establish ongoing eligibility by Standard Score or Informed Clinical Opinion.

For more information, please refer to Baby Watch's Eligibility Criteria policy.